arXiv:1610.04389v1 [hep-ex] 14 Oct 2016

Frascati Physics Series Vol. 64 (2016)
Frontier Objects in Astrophysics and Particle Physics
May 22-28, 2016

NEW PROJECTS ON DARK PHOTON SEARCH

Venelin Kozhuharov
University of Sofia, 5 J. Bourchier Blvd, 1164 Sofia, Bulgaria
INFN - LNF, Via E. Fermi 40, 00044 Frascati (RM), Italy
Mauro Raggi
INFN - LNF, Via E. Fermi 40, 00044 Frascati (RM), Italy
Paolo Valente
INFN - Sezione di Roma, P.le Aldo Moro, 2 - 00185 Roma - Italy

Abstract

Despite the great success of the Standard Model of particle physics the nature of Dark Matter still remains unclear. Recently, the idea of the existence of a hidden sector coupling only weakly with the ordinary matter was revitalized and gained popularity. A simple mediator between the hidden and the visible sector could be a vector particle of a new gauge interaction, the so called dark photon. Numerous of activities were initiated to probe its parameter space. The present results and the foreseen experiments aimed to search for dark photons in few directions are reviewed and discussed.

1 Introduction

The Standard Model (SM) of particle physics provides a unique description of almost all phenomena in the microworld. However, the unexplained nature

of Dark Matter indicates the necessity of its extension to a more fundamental theory, which incorporates also the cosmological observations. This can be realized in a large variety of theoretical models.

From experimental point of view the best path to follow is to focus on few of the observed "smoking guns" indicating a possible discrepancy between the SM prediction and the results from the data. At present there are still unexplained anomalies with the annually modulated excess of signal observed by DAMA/Libra [1]; the positron and antiproton excess in the cosmic rays [2, 3, 4, 5]; the three sigma discrepancy in the anomalous magnetic moment of the muon [7]; the failure to explain the ^{8}Be anomaly [6] within nuclear physics. All those measurements may indicate the presence of new undiscovered so far particles.

An elegant model to address the possible existence of new degrees of freedom is the concept of a dark sector (DS) of particles interacting only weakly with the Standard Model particles. The origin of the interactions could be due to the presence of a mediator - an object carrying both SM and DS quantum numbers or SM fields possessing a relatively small charge under any of the gauge symmetries in the Dark Sector. In that sense the mediator provides a so-called portal to the DS [8] [9]. One of the simplest possible realization that is ultra violet safe (i.e. does not introduce new scale in the lagrangian) is by employing a vector gauge field, the so-called Dark Photon A' (DP), which interacts weakly with the SM fermions

$$\mathcal{L} \sim g' q_f \bar{\psi}_f \gamma^\mu \psi_f A'_\mu, \tag{1}$$

where g' is the universal coupling constant and q_f are the corresponding fermion charges. The term in equation (1) could be effectively realized also through kinetic mixing of the massive DP with the ordinary photon. In this picture the interaction of the SM particles with the dark photon will be described by two parameters - $\epsilon \sim g' q_f$ and the dark photon mass $m_{A'}$. The coupling parameter ϵ could be flavour dependent giving rise to different leptophobic or leptophilic DP models [10].

Two different scenarios depending on the phenomenology in the dark sector could be identified. In the case when no new light degree of freedom χ exists in the dark sector the dark photon will decay to SM particles only, with $\mathrm{Br}(A' \to e^+e^-) = 100\%$ for 1 MeV $< m_{A'} <$ 210 MeV. If, however, $m_\chi < m_{A'}/2$ then the decay $A' \to \chi\chi$ will be dominant since it is not suppressed

by the small value of ϵ. In the latter scenario the observables will depend on two additional parameters, the coupling strength in the dark sector α_D and m_χ. These two scenarios, so called "visible" and "invisible", result in very different experimental signature. In the visible case a narrow resonant might be observed in the dilepton or, in general, in the di-particle invariant mass spectrum, while in the invisible case the existence of a DP could present itself through the search for missing mass or missing energy.

2 Visible dark photon decays

Currently, most of the experiments addressing the existence of an A' performed "peak" searches in the e^+e^- invariant mass spectrum. This approach requires a precise spectrometer providing measurement of the electron momentum with high precision. The production of the A' could be either through a bremsstrahlung process (in electron-on-target experiments) or through the $e^+e^- \to \gamma A'$ at electron-positron colliders.

Two new dedicated to the A' search experiments are planed in the near future - the HPS experiment at the JLaB [12, 11] and the MAGIX experiment at Mainz [13]. Both of them will exploit an electron beam impinging on a thin target.

The HPS experiment will measure the momentum of A' products using a silicon vertex tracker placed inside a dipole magnet. A downstream lead tungstate calorimeter will serve for fast energy measurement and triggering. The silicon tracker is made of six dual sensor layers and will allow to address the dark photon decays in two different regions of its parameter space. For large values of ϵ^2 ($\epsilon^2 > 10^{-7}$) A' decays promptly and the event selection will be based on the e^+e^- invariant mass reconstruction. The dominant background originates from the internal pair conversion of the bremsstrahlung photon into e^+e^- pair. When $\epsilon^2 \leq 10^{-8}$ the finite lifetime of A' will result in the reconstruction of vertices displaced from the interaction point inside the target. This channel is particularly interesting since it could be background free, at the expense of a low signal yield. The projected sensitivity for HPS is $\epsilon^2 < 3-4\times10^{-7}$ for 20 MeV $\leq m_{A'} \leq$ 300 MeV and $2\times10^{-8} \leq \epsilon^2 \leq 2\times10^{-10}$ in the region 20 MeV $\leq m_{A'} \leq$ 200 MeV

The MAGIX experiment is planned to operate at the new energy-recovering superconducting accelerator at Mainz, MESA, which provides an electron beam

with energy up to 155 MeV and 1 mA beam current. The target will be accomplished as a windowless supersonic gas jet with high density ($10^{19}/cm^2$). The A' decay products, e^+ and e^-, will be detected in a double arm spectrometer with planned resolution of $\delta p/p = 10^{-4}$. The projected sensitivity indicates reach of $\epsilon^2 \sim 10^{-8}$ for masses 10 MeV $\leq m_{A'} \leq$ 50 MeV.

3 Invisible dark photon decays

The mass spectrum of the particles in the dark sector is in general ambiguous and nothing prevents the existence of light states. Such a scenario is relatively difficult to probe due to the impossibility to perform a complete reconstruction of the final state when A' decays to $\chi\chi$. Thus it is important to start with an initial state that can be fully described. The annihilation of e^+ with e^- is one of the possibility while another is to usage of mono-energetic beam and to search for missing energy taken away by the produced dark photons.

3.1 Missing mass technique

The missing mass technique relies on the complete reconstruction of the annihilation process

$$e^+ + e^- \rightarrow \gamma A' \tag{2}$$

through the measurement of the energy and the direction of the recoil photon. In positron on target collisions the missing mass squared is then computed as

$$M^2_{miss} = (P_e + P_{beam} - P_\gamma)^2 \tag{3}$$

where usually the electron is considered to be at rest ($P_e = (m_e, 0, 0, 0)$). The cross section for the production of A' is enhanced with respect to the ordinary $e^+e^- \rightarrow 2\gamma$ process by a factor δ [14], especially for $m_{A'}$ close to the centre of mass energy of the interaction. The dominant background processes are listed in table 1 and originate from the bremsstrahlung emission in the field of the target nuclei and from the $e^+ + e^- \rightarrow \gamma\gamma\gamma$ annihilation. Usually, the measurement of the energy of the recoil photon provides missing mass squared resolution good enough to suppress the $e^+ + e^- \rightarrow \gamma\gamma$ background.

Three experiments, PADME at the DAΦNE Linac at LNF-INFN [15], MMAPS at Cornell [18], and at VEPP3 at Novosibirsks [17], are planning to exploit this technique. They share similar design properties.

Table 1: *Dominant background contributions to the missing mass technique*

Background process	σ ($E_{beam} = 550$ MeV)	Comment
$e^+e^- \to \gamma\gamma$	1.55 mb	
$e^+N \to e^+N\gamma$	4000 mb	$E_\gamma > 1MeV$, on carbon
$e^+e^- \to \gamma\gamma\gamma$	0.16 mb	$E_\gamma > 1MeV$, CalcHEP [16]
$e^+e^- \to e^+e^-\gamma$	188 mb	$E_\gamma > 1MeV$, CalcHEP

PADME experiment at LNF-INFN, shown in fig. 1, will use 550 MeV positron beam impinging on a 100 μm thick active target which is made of polycrystalline diamond. The recoil photons from the process $e^+ + e^- \to \gamma A'$ will be detected by a ring-shaped BGO crystal calorimeter, which is located 3 m downstream and provides energy and position information. The non-interacted positron beam will be deflected outside the acceptance of the calorimeter by a dipole magnet. Three sets of plastic scintillator detectors will serve to register the charged particles and will provide an efficient veto for the bremsstrahlung background. In addition, a fast detector for photons, placed along the undeflected beam axis, will help with the suppression of the three photon annihilation background. The complete setup is located in vacuum to diminish the possible beam-residual gas interactions.

The approach of VEPP3 is to use an internal hydrogen gas target placed at the Novosibirsks storage ring, illuminated by the 500 MeV e^+ beam. Both the VEPP3 and the MMAPS experiments aim to use CsI crystals from CLEO for their calorimeters, which are shown to provide $\sigma_E/E = 3\%$ for 180 MeV positrons. In order to be able to operate in parallel with the ongoing VEPP3 activities an extension of the existing beam line is proposed. This would allow to place a charged particle veto detector to suppress the background. MMAPS uses a beryllium target and will profit from the higher beam energy - up to 5.3 GeV. This extends MMAPS sensitivity region to masses up to $m_{A'} \sim 74$ MeV.

A comparison of the characteristics and the performance of the PADME, MMAPS, and VEPP3 experiments is shown in table 2. PADME will be the first experiment to run and possible upgrades of the DAΦNE linac could extend its sensitivity down to the region of $\epsilon^2 \sim 3 - 5 \times 10^{-7}$. The VEPP3 experiment requires an approval of beam line modification while the Cornell MMAPS experiment is in process of identification of finding.



Figure 1: *CAD schematics of the PADME experiment at Frascati Linac.*

3.2 Missing energy technique

During the shower development inside a calorimeter, the dark photons could be created through an A'-strahlung process by any of the secondary particles. They could account for a large part of the undetected energy of the primary particle and their existence may manifest itself in the form of a large fraction of events with missing energy. This technique relies on the precise knowledge of the response of a calorimeter to an electromagnetic shower development. It is exploited by the recently approved NA64 experiment operating at CERN SPS [19]. Synchrotron radiation tagged 100 GeV e^- beam with momentum determined in a magnetic spectrometer impinges on an electromagnetic calorimeter (ECAL) acting as an active beam dump. Events with less than 50 GeV energy in the ECAL and no signal in the veto and in the downstream hadron calorimeter are considered as signal. In the hypothesis of zero background observation the NA64 experiment aims to set limits down to $\epsilon^2 \sim 10^{-8} - 10^{-6}$ for A' masses in the region 10 MeV - 100 MeV, assuming 10^{10} electron events.

3.3 Scattering of dark matter particles

Another possible indication of the presence of dark photons decaying into invisible particles χ could be the direct observation of the χ scattering in a low

Table 2: *Comparison between the experiments exploiting the missing mass technique*

	PADME	MMAPS	VEPP3
Place	LNF	Cornell	Novosibirsk
Beam energy	550 MeV	Up to 5.3 GeV	500 MeV
$M_{A'}$ limit	23 MeV	74 MeV	22 MeV
Target thickness [e^-/cm^2]	2×10^{22}	$O(2 \times 10^{23})$	5×10^{15}
Beam intensity	8×10^{-11} mA	2.3×10^{-6} mA	30 mA
$e^+e^- \to \gamma\gamma$ rate [s^{-1}]	15	2.2×10^6	1.5×10^6
ϵ^2 limit (plateau)	10^{-6}	10^{-6} - 10^{-7}	10^{-7}
Time scale	2017-2018	?	2020 (ByPass)
Status	Approved	Not funded	Proposal

noise detector. This approach is similar to the beam dump technique widely used in the past to search for milicharged particles [9]. The A' are produced, through A'-strahlung and then decay to $\chi\chi$ pair. The BDX experiment [20], proposed to take place at JLaB, will use a CsI (Tl) detector shielded by an active vetoing system to suppress the cosmogenic background. Studies indicated that sensitivity down to $\epsilon^2 \sim 10^{-9}$ for $m_{A'} = 50$ MeV could be reached. However, this depends on the extra parameters m_χ and α_D.

4 Conclusions

Present searches for new vector particle A' aim to cover the visible and the invisible final state scenarios in parallel. While on the visible side many results appeared in the last decade the parameter space for A' decaying predominantly to non Standard Model particles is being addressed just recently. The new activities in this direction can be summarized in fig. 2. The PADME and the NA64 experiments are approved and are expected to provide interesting results by 2020. At the same time few other unique projects are in preparation.

5 Acknowledgements

The authors would like to thank the organizers for the invitation to present at the Vulcano workshop. VK acknowledges support from INFN, LNF-INFN under contract LNF-SU 70-06-497/07-10-2014 and University of Sofia science



Figure 2: *Recent perspectives for invisible dark photon searches.*

fund under grant FNI-SU N39/2016.

References

1. R. Bernabei *et al.*, Eur. Phys. J. **C67**, 39 (2010).
2. O. Adriani *et al.* [PAMELA Collaboration], Nature **458**, 607 (2009).
3. M. Ackermann *et al.* [Fermi LAT Collaboration], Phys. Rev. Lett. **108**, 011103 (2012).
4. M. Aguilar *et al.* [AMS Collaboration], Phys. Rev. Lett. **110**, 141102 (2013).
5. AMS-02 Collaboration, Talks at the "AMS Days at CERN", 15-17 april 2015
6. A.J. Krasznahorkay *et al.*, Phys. Rev. Lett. **116** (2016) 042501.

7. G. W. Benett *et al.* (The g-2 Collaboration), Phys. Rev. **D73** (2006) 072003.

8. R. Essig *et al.*, (2013), arXiv:1311.0029 [hep-ph].

9. T. Spadaro, these proceedings.

10. M. Raggi and V. Kozhuharov, Riv. Nuovo Cim. 38, no. 10, 449 (2015).

11. T. Nelson, talk at "Dark Sectors Workshop", 28-30 Apr., SLAC (2016).

12. A. Celentano [HPS Collaboration], J. Phys. Conf. Ser. **556**, no. 1, 012064 (2014)

13. A. Denig, talk at "Dark Sectors Workshop", 28-30 Apr., SLAC (2016).

14. M. Raggi and V. Kozhuharov, Adv. High Energy Phys. **2014**, 959802 (2014)

15. M. Raggi, V. Kozhuharov, P. Valente et al., The PADME experiment Technical Proposal.

16. A.Pukhov *et al.*, Preprint INP MSU 98-41/542,arXiv:hep-ph/9908288
A.Pukhov, e-Print Archive: hep-ph/0412191.

17. B. Wojtsekhowski, D. Nikolenko and I. Rachek, arXiv:1207.5089 [hep-ex]., also talk at "Dark Sectors Workshop", 28-30 Apr., SLAC (2016).

18. J. Alexander, talk at "Dark Sectors Workshop", 28-30 Apr., SLAC (2016).

19. S. Andreas *et al.*, arXiv:1312.3309 [hep-ex].

20. M. Battaglieri *et al.* [BDX Collaboration], arXiv:1406.3028
M. Battaglieri *et al.* [BDX Collaboration], arXiv:1607.01390 [hep-ex].